- 26. (canceled)
- 27. The method of claim 24, wherein said oxidized amino acid residue is histidine.
- 28. The method of claim 24, wherein said oxidized amino acid residue is tryptophan.
- 29. The method of claim 24, wherein said aflibercept variant is selected from an amino acid residue on a polypeptide having an amino acid sequence as set forth in the group consisting of: SEQ ID NO.: 17, SEQ ID NO.: 18, SEQ ID NO.: 19, SEQ ID NO.: 20, SEQ ID NO.: 21, SEQ ID NO.: 22, SEQ ID NO.: 23, SEQ ID NO.: 56, SEQ ID NO.: 64, SEQ ID NO.: 65, SEQ ID NO.: 66, SEQ ID NO.: 67, SEQ ID NO.: 68, SEQ ID NO.: 69, SEQ ID NO.: 70, SEQ ID NO.: 71, and combinations thereof.
- 30. The method of claim 24, further subjecting said protein sample of step (b) to one or more additional chromatographic steps selected from the group consisting of:

- cation exchange chromatography, hydrophobic interactive chromatography, size exclusion chromatography and a combination thereof.
- 31. The method of claim 19, wherein said oxidized amino acid residue is histidine.
- 32. The method of claim 19, wherein said oxidized amino acid residue is tryptophan.
- **33**. The method of claim **19**, wherein said cell is selected from the group consisting of CHO, NS0, Sp2/0, embryonic kidney cells and BHK.
- 34. The method of claim 19, further subjecting said protein sample of step (b) to one or more additional chromatographic steps selected from the group consisting of: cation exchange chromatography (CEX), hydrophobic interactive chromatography, size exclusion chromatography and a combination thereof.

\* \* \* \* \*